

## Examining the variations of ground level ozone and nitrogen dioxide in a rural area influenced by brick kiln industries

K. Elampari, T. Chithambarathanu and R. Krishna Sharma  
Research Department of Physics, S.T. Hindu College, Nagercoil, India- 629002  
elampari@rediffmail.com

### Abstract

Burning of biomass fuels is responsible for the emission of both trace and non-trace greenhouse gases such as CO<sub>2</sub>, CH<sub>4</sub>, CO, N<sub>2</sub>O, NO<sub>x</sub> and NO. The photolysis of nitrogen dioxide (NO<sub>2</sub>) directly leads to the formation of ozone (O<sub>3</sub>) in the lower atmosphere. This work is an attempt to estimate the level of ground level ozone (GLO) and NO<sub>2</sub> concentrations in the atmospheric surface layer of a rural area dominated by traditional brick kiln industries. A portable monitor with O<sub>3</sub> and NO<sub>2</sub> sensor heads was employed for the measurement of the trace gases from May 2009 to July 2009. The GLO concentration showed a clear diurnal cycle with higher values in the daytime and notably lower values at night time. The concentration of GLO during the period of study varied from a minimum value of 7 ppb around 0530 h to a value of 48 ppb around 1430 h. The lowest mean day time value of NO<sub>2</sub> was 2.52 ppb and the highest mean night time value was 5.83 ppb.

**Keywords:** Brick kilns, biomass, diurnal variation, ozone precursor, photochemical reaction, ground level ozone

### Introduction

Ozone is an important trace species present in the atmosphere. It is produced photochemically in the lower atmosphere from reactions involving a variety of volatile organic compounds (VOCs) composed mainly of non-methane hydrocarbons in the presence of sufficient sunlight and nitrogen oxides (NO<sub>x</sub>) such as nitric oxide (NO) and nitrogen dioxide (NO<sub>2</sub>). NO<sub>x</sub> are emitted from a variety of natural and anthropogenic sources (Al-Khalaf, 2006). Ozone is the most important gas in the photochemistry of the atmosphere and is the primary constituent of photo-oxidative smog. Thus, it can be considered as an indicator of the overall burden of the atmospheric oxidants. The physical and chemical processes of this GLO directly depend on the local precursor emission levels and meteorological factors. In the ambient air even a micro level variation in the concentration of GLO brings lethal effects on the biological and material systems (Ravishankar & Rao, 2002; Moller, 2004). Ozone in the troposphere has been steadily increasing at the rate of 0.5-2% per year (Vingarzan, 2004) over the past few decades. Thus, abundance of GLO concentration is increasing at many sites over the globe and expected to rise significantly above the threshold limit set for human (hourly average of 80 ppbv set by World Health Organization) throughout the 21<sup>st</sup> century (WHO, 2000; IPCC, 2001). Further increase of tropospheric ozone is anticipated under likely scenarios of increasing regional and global emissions of ozone precursor compounds (Seinfeld, 1991). The two sources of ozone in the natural troposphere are: a) Injected from stratosphere and b) Photochemical production via NO<sub>x</sub>-VOC reactions in sunlight involving naturally occurring oxides of nitrogen and hydrocarbons or CO (Seinfeld & Pandis, 2006; Salve *et al.*, 2007). Studies show that apart from the industrial and vehicular exhausts, fumes from brick kilns also contribute to the

increase in the level of local ozone at surface levels (Pudasainee *et al.*, 2006). Using biomass fuels in traditional brick industries is responsible for the emission of both trace and non-trace gases such as CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, NO<sub>x</sub> and NO and also toxic fumes containing suspended particulate matters rich in carbon particles and high concentration of carbon monoxides and oxides of sulphur (SO<sub>x</sub>) (Alam, 2006). Therefore, brick making industries can be considered as one of the important sources of greenhouse gases.

The objective of this study is to determine the level of GLO and its precursor NO<sub>2</sub> concentrations in a rural area where a large number of traditional brick kilns are active.

### Study area

The study area Chenbagamanputhur (8°15'1" N, 77°29'19" E) is a rural place in Kanyakumari district and is about 12 km from Nagercoil town (Fig. 1). Brick production is one of the traditional crafts of the people in this village and is a valuable source of employment. There are approximately 300 conventional brick kilns around this study place. These kilns are heavily utilized during dry season since the wet weather in other seasons prevents the bricks from setting hard. In this brick kilns, firewood is used as the main fuel and approximately 1,000 bricks need 45 kg of fire wood.

### Materials and methods

Ground level ozone concentration and nitrogen dioxide measurements were carried out in the rural site Chenbagamanputhur. A portable Aeroqual series S200 ozone monitor similar to that of Aeroqual S300 used by Dovile (2009) was used for the measurement. An Aeroqual series 200 ozone monitor is constructed to measure low and high ozone levels. Its ultra low concentration ozone head measures the ozone concentration from 0.000 to 0.500 ppm and a high

concentration ozone head measures the ozone concentration from 0.50 to 20.00 ppm. Accuracy of a low concentration ozone head is +/- 0.010 ppm (from 0 to 0.100 ppm); +/- 10% (0.100 to 0.500 ppm), while that of a high concentration ozone head is +/- 10% (from 0.20 to 2.00 ppm); +/- 15% (from 2.00 to 20.00 ppm), the measurement units being either ppm or  $\mu\text{g}/\text{m}^3$ . The operating temperature range is from 5°C to 50°C, relative humidity limits are 5% and 95%. **Similar kind of NO<sub>2</sub> sensor has been used for nitrogen dioxide measurement. A gas sensitive semiconductor (GSS) type sensor is described in [www.aeroqual.com](http://www.aeroqual.com).** Sampling was carried out for three months from May 2009 to July 2009. For ozone seven readings were taken per day starting from 0530 h to 2330 h with 3 h interval. For NO<sub>2</sub>, only two readings were taken one at day time (1430 h) and the other at night time (2330 h).

### Results and discussion

The results of surface ozone and NO<sub>2</sub> levels observed during the present study are given below.

#### Diurnal variation of GLO

The observed mean values of GLO and nitrogen dioxide are shown in Table 1 & 2. The GLO concentration showed a clear diurnal cycle with higher values in the daytime and notably lower values at night time. This diurnal pattern is shown in Fig. 2. It can be seen that the ozone concentration was low at the early morning hours (0530 h), it slowly picked up in the forenoon and then built up faster and ultimately reached the maximum value in the afternoon. A gradual decrease was observed in the evening and a sudden fall was recorded after the sunset.

Table 1. Monthly mean GLO concentration (in ppb)

Month	Time						
	5:30	8:30	11:30	14:30	17:30	20:30	23:30
May 2009	20.40	26.40	32.40	44.60	33.60	30.20	27.40
June 2009	10.00	15.17	23.67	31.33	28.00	20.67	18.00
July 2009	19.33	23.83	30.00	36.67	34.33	27.83	23.67

The diurnal variation of GLO is helpful to understand the different processes responsible for ozone formation and destruction at this particular location. Chemical and atmospheric dynamic processes regulate the diurnal ozone concentration. The maximum value of GLO was observed around post noon which indicates that there exists a direct coincidence of GLO variation with the surface temperature. The photochemical production of ozone depends on the concentration of its precursor gases and meteorological parameters. The variation in these parameters changes the diurnal magnitude of ozone concentration. The lower levels of GLO observed during night time may be due to chemical loss of ozone (NO<sub>x</sub> titration) and also due to the ozone loss at the surface of the earth, as there is no production of ozone during night hours (Debaje *et al.*, 2003). The concentration of GLO during the period of study varied from a minimum value of 7 ppb around 0530 h to a value of 48 ppb around 1430 h



Research article

©Indian Society for Education and Environment (ISEE)

whereas the monthly mean concentration of GLO varied from 10 ppb to 44.60 ppb. May 2009 recorded the maximum and June 2009 recorded the minimum values of GLO.

#### Behaviour of NO<sub>2</sub>


The NO<sub>x</sub> emission from the brick kilns has a major role in the formation of ozone. The presence of NO<sub>x</sub> is very important because NO<sub>2</sub> can be the only potential source of ozone in the study area. If there is a fractional amount of NO<sub>x</sub> in the troposphere, it receives solar radiation and the ozone is formed as indicated by the following reactions in eq. 1 and 2 (Tang, 2009).



Where, O (<sup>3</sup>P) represents the oxygen atom in the ground state. However if those are the only reactions, then the ozone concentration would attain a steady state and does not reach to high value because the reaction (3) will occur in which ozone will be destroyed by NO produced in the reaction (1).



But with the presence of any hydrocarbons in the atmosphere, NO will be transformed to NO<sub>2</sub> by peroxy radicals such as RO<sub>2</sub><sup>\*</sup> and hydroxyl-radicals HO<sub>2</sub><sup>\*</sup> as indicated in eq. (4) and (5) (Tang, 2009).



The RO<sub>2</sub><sup>\*</sup> and HO<sub>2</sub><sup>\*</sup> radicals originate in the oxidation of the VOCs. Hence no ozone is consumed for the NO oxidation and the concentration of O<sub>3</sub> is no longer constant, but increases. NO<sub>2</sub> produced in the above reactions is then used to form ozone through reactions of eq. (1) and (2). As a result the concentration of ozone becomes high under the condition of strong solar radiation (Seinfeld & Pandis, 2006).

The observed levels of NO<sub>2</sub> concentration is shown in Fig. 3. The mean concentration of NO<sub>2</sub> at night was greater than that of day (i.e.,) 5.83, 4.56 and 4.77 ppb at night and 2.79, 2.64 and 2.52 ppb at day during the study period. This behavior of NO<sub>2</sub> indicates that the diurnal variation of ozone depend on photochemical reactions during day time. The low NO<sub>2</sub> concentration observed during day time may be due to the photochemical reaction taking place for ozone production using NO<sub>2</sub> as a fuel. The maximum of GLO at noon is related to the highest photolysis rate of NO<sub>2</sub> at noon.

Thus NO<sub>2</sub> acts as a precursor of ozone. After sunset, the intensity of photochemical reaction reduces and hence NO<sub>2</sub> concentration builds up. Table 3 shows the observed minimum and maximum values of GLO and NO<sub>2</sub> during the study.

#### Comparison of GLO & NO<sub>2</sub> concentrations

Analyzing the comparative diagram (Fig. 4) of GLO and NO<sub>2</sub> concentration the relationship between the two trace species in a monthly scale could be obtained. The two gases, one as a precursor (NO<sub>2</sub>) and

Table 2. Monthly mean NO<sub>2</sub> concentration (in ppb)

Month	Day	Night
May 2009	2.79	5.83
June 2009	2.64	4.56
July 2009	2.52	4.77

"Ozone"

<http://www.indjst.org>

Elampari *et al.*  
Indian J.Sci.Technol.

Fig. 1. Location map of the observation site




Fig. 2. Mean GLO concentration for May, June & July 2009.




Fig. 3. Mean NO<sub>2</sub> concentration in May, June & July 2009





Fig. 4. Variation of GLO and NO<sub>2</sub> from May to July 2009



the other as a product (GLO) vary in a similar way but with a different range of values throughout the period of observation. It clearly indicates that the role of photochemical process and emission rates at this observational location are significant. One more interesting inference from the fig. 4 is that the increasing or decreasing trend of NO<sub>2</sub> started first and then followed by ozone with the delay of at least a week's time. This is also helpful in substantiating the idea of NO<sub>2</sub> acting as a precursor emitted from both natural and man-made sources, for the photo chemical production of ozone (Ahammed *et al.*, 2006).

Table 3. The maximum and minimum concentrations of GLO & NO<sub>2</sub> (in ppb)

Month	GLO		NO <sub>2</sub>	
	Minimum	Maximum	Minimum	Maximum
May 2009	18.00	48.00	2.35	7.25
June 2009	7.00	34.00	1.75	6.11
July 2009	10.00	40.00	1.26	7.47

The local meteorological factors and the activities involved in the brick kilns played a great impact on the observed pattern. In May 2009 with clear sky conditions and intense solar flux density marking the peak summer season at this place which provided a conducive atmosphere for the better production of NO<sub>2</sub> and hence ozone. June 2009 mostly had cloudy and rainy days. Hence lower concentrations of GLO and NO<sub>2</sub> were recorded than that of July 2009 and a similar condition was observed by Muralidharan *et al.* (1989). The recovery of background values of these gas concentrations in July 2009 may be due to the reversal of meteorological conditions from the previous month.

### Conclusion

As a case study at Chenbagaramanputhur village, the ambient concentration of GLO and NO<sub>2</sub> and the relationship between these two generic species of gases has been revealed. Due to the improper construction of kilns large amount of fumes were released which may contain gases like CO, CO<sub>2</sub> and NO<sub>x</sub>. Hence these brick kilns are acting as a point source of the precursor gases of ozone. During May, which is the crucial period for kiln works, large amount of fumes were released from kilns and hence maximum concentration of GLO was observed. Because of rain during June the activities were reduced and hence the values of GLO were comparatively lower than the other two months. Once again work was regained in July, which resulted into improved emissions and increased concentrations. The GLO and NO<sub>2</sub> levels were within the permissible limits of both national and international air quality standards. The monthly average values were found to be explicitly depended on natural and anthropogenic emissions as well as the photochemical mechanism.

### Acknowledgement

We thank the Principal, Chairman-Secretary, S.T. Hindu College, Nagercoil for providing the facilities to carry out this study.

Research article

©Indian Society for Education and Environment (ISEE)

### References

- Ahammed YN, Reddy R and Rama Gobbal RK (2006) Seasonal variation of the surface ozone and its precursor gases during 2001-2003, measured at Anantapur, a semi arid site in India. *Atmos. Res.* 80(2-3), 151-164.
- Alam SA (2006) Use of biomass fuels in the brick-making industries of Sudan: implications for deforestation and greenhouse gas emission, Dept. of forest ecology, Univ. of Helsinki, Finland. pp:16-17.
- Al-Khalaf AK (2006) Influence of meteorological and related factors on surface ozone pattern at Makkah station. *J. Environ. Sci.* 11,1-19
- Debaje SB, Johnson S Jeyakumar, Ganesan G and Jadhav DB (2003) Surface ozone measurements at tropical rural coastal station Tranquebar, India. *Atmos. Environ.* 37, 4911-4916.
- Intergovernmental Panel on Climate Change (IPCC) (2010) Electronic reference. Retrieved Jan. 1, 2010, from [http://www.grida.no/climate/ipcc\\_tar/wg1/index.htm](http://www.grida.no/climate/ipcc_tar/wg1/index.htm)
- Moller D (2004) The tropospheric ozone problem. *Arh Hig Rada Toksikol.* 55(1), 11-25.
- Muralidharan V, Mohan Kumar G and Sampath S (1989) Surface ozone variation with rainfall. *Earth Environ. Sci.* 130, 1.
- Pudasainee D, Sapkota B, Shrestha ML, Kaga A, Kondo A and Inoue Y (2006) Ground level ozone concentrations and its association with NO<sub>x</sub> and meteorological parameters in Kathmandu valley, Nepal. *Atmos. Environ.* 40, 8081-8087.
- Ravi Shankar P and Rama Rao G (2002) Impact of air quality on human health: A case study of Mumbai City, India, IUSSP Regional Conf. on Southeast Asia's Population in a Changing Asian Context.
- Salve PR, Satapathy DR, Katpatal YB and Wate SR (2007) Accessing spatial occurrence of ground level ozone around coal mining areas of Chandrapur district, Maharashtra, India. *Environ. Monit. Assess.* 133, 1-3.
- Seinfeld JH (1991) Rethinking the Ozone problem in urban and regional air pollution. National Acad. Press, Washington, DC
- Seinfeld JH and Pandis SN (2006) Atmospheric physics and chemistry from air pollution to climate change. John Wiley & Sons, Inc. NY. pp:234-336.
- Tang L (2009) Regional and local surface ozone variations in relation to meteorological conditions in Sweden. Dept. of Earth Sci., Univ. Gothenburg.
- Vingarzan R (2004) A review of surface ozone background levels and trends. *Atmos. Environ.* 38, 3431-3442.
- World Health Organisation (WHO) (2009) Electronic reference. Retrieved Nov.10, 2009 from <http://www.airimpacts.org/documents/local/AQGUIDE.pdf>.